

“Consistency and hard work  
are key to success”

# CSIR NET – Life Science





## Unit 1: Biochemistry

13

Enzyme Regulation  
and Metabolism



Order Online and Get  
Free Delivery Across India





## Points to be covered in this Lecture



Allosteric Enzyme



Isoenzyme



Ribozyme



Regulation of Enzyme Activity



Enzyme Purification



Basics of metabolism



Glycolysis



## Allosteric Enzymes

Intracellular ✓


Pacemaker ← Slow enzyme

Catalytic + Regulatory subunit ✓

Intrinsic regulation ← Effectors

Cooperativity → multiple subunit



Hill's equation → Donot obey mm equation





## Hill's Equation and Plot

Describe the cooperative binding of substrate to allosteric enzyme



$$V_0 = V_{max} \frac{[S]^n}{K_{0.5} + [S]^n}$$



$n$  = Hill's coefficient  
 $n$  value is more than 1

$$K_m = K_{0.5}$$

$[S]$  required to achieve  $V_0 = \frac{1}{2} V_{max}$

$$V_{max} = K_{cat} E_T$$

## Cooperativity






Straight Line

$$V_0 = V_{max} \frac{[S]^n}{K_{0.5} + [S]^n}$$

$$\frac{\log V_0}{V_{max} - V_0} = n \log [S] - \log K_{0.5}$$

$$Y = m \cdot X - (C)$$



X axis =  $\log [S]$

Y-axis =  $\frac{\log V_0}{V_{max} - V_0}$

X-axis intercept =  $\log K_{0.5}$

Slope = n (Hill's coefficient)



## Allosteric constant (L)


The equilibrium constant for the transition between two forms of an allosteric protein in the absence of regulator

$$L = T_0 / R_0$$

$$L = \frac{\text{Tense state} \leftarrow \text{Inhibitor } (-)}{\text{Relax state} \leftarrow \text{Activators } (+)}$$

Positive regulator: Decreases allosteric constant

Negative regulator: Increases allosteric constant






## I. Isozymes or isoenzymes:

- Variants of the enzyme that exist co-functionally within the same organism
- Arise from different genes or from alternative splicing of the same gene.
- Differ in terms of molecular weight, optimal pH, kinetic parameters like  $K_m$  and  $K_{cat}$  and regulatory mechanisms.
- Often expressed in specific tissues or at certain developmental stages.
- Allow for more complex regulation of metabolic pathways
- Can be markers for tissue damage or disease states






### Lactate dehydrogenase (LDH)



- LDH is composed of four subunits, which can be of two types: H (heart) and M (muscle).
- The different combinations of these subunits form five isoenzymes:


- ✓ LDH-1 (H<sub>4</sub>): heart and red blood cells,
- ✓ LDH-2 (H<sub>3</sub>M<sub>1</sub>): reticuloendothelial system,
- ✓ LDH-3 (H<sub>2</sub>M<sub>2</sub>): in lungs,
- ✓ LDH-4 (H<sub>1</sub>M<sub>3</sub>): in kidneys, pancreas, and placenta,
- ✓ LDH-5 (M<sub>4</sub>): liver and skeletal muscle.








## Creatine Kinase distribution



## Creatine Kinase



- ✓ Dimer
- ✓ Two types of subunit M (Muscle) and B (Brain)
- CK has three isoenzymes
  - CK-MM**: Found predominantly in skeletal muscle and heart.
  - CK-MB**: Found in heart muscle and is of particular importance in diagnosing myocardial infarction.
  - CK-BB**: Primarily present in the brain and smooth muscle, including the gastrointestinal tract.



Ribo-enzyme → Ribonucleic acid (RNA) act as enzyme

- Thomas Cech and Sidney Altman
- Catalytic RNA
- Ancestral enzymes *Limited conformations*
- Slow and Unstable, *Limited catalytic strategy (2'-OH)*

✓ **Hammerhead ribozyme:** Catalyzes its own cleavage during replication of viroids and virusoids.

✓ **Hairpin ribozyme:** Also involved in self-cleavage linked to replication processes.

*Group I = Pre rRNA*


*Group II : Pre mRNA*

✓ **Group I and Group II introns:** Catalyze their own excision from RNA transcripts and the ligation of the remaining exons.

✓ **RNase P:** Involved in the maturation of pre-tRNA by cleaving its 5' leader sequence.

- **Peptidyl transferase of the ribosome:** forms peptide bonds

↳ 23 S rRNA (Prokaryotes)







## REGULATION OF ENZYME ACTIVITY

### I. Allosteric regulation : Allosteric enzyme

✱ Homotropic effectors: When substrate itself act as effector

Heterotropic effectors: Except substrate some other molecule act as effector


{  
 Binds to  
 Allosteric site  
 {  
 Non-covalent  
 interaction  
 (Reversible)






## Feedback inhibition

Type of negative allosteric regulation where the end product of a metabolic pathway inhibits an enzyme involved earlier in the pathway.



## Feed forward activation


Type of **positive** allosteric regulation where a **metabolite produced early** in a pathway enhances the activity of an enzyme later in the same pathway.



## 2. Regulation of enzymes by covalent modification


Reversible addition or removal of chemical groups to specific amino acids in an enzyme, thereby altering its activity, stability, or cellular location

- Phosphorylation/Dephosphorylation: ser, Tyr, His, Asp, Thr  
Kinase      Phosphatase ← Regulatory enzyme.




• Nucleotidylation of a specific Tyr : Glutamine synthetase (E. coli).

- Addition of nucleotide - ADP
- Enzyme is inactivated



• ADP-ribosylation of a specific Arg : Adenylate cyclase of E. coli


- Addition of ADP ribose from NAD
- enzyme is inactivated



N = Nicotinamide

• Disulphide reduction

- Reducing env<sup>y</sup>
- Disulphide bond break
- Sulphydryl bond are formed
- enzyme is activated






### 3. Proteolytic activation — Irreversible — Enzyme has short half life

- The precise cleavage of one or more specific peptide bonds within a zymogen.
- This cleavage induces a conformational change in the molecule, either by removing a segment that inhibits activity or by exposing a previously hidden active site.
- Cascading Events

#### Examples

- Digestive enzymes are secreted as zymogens (pepsinogen, trypsinogen, chymotrypsinogen).
- The coagulation cascade (blood clotting)
- Apoptotic cascade
- Complement protein cascade









#### 4. Substrate Inhibition:

- Enzyme has more than one binding site for the substrate.
- While the first binding site is catalytic, additional binding at a second regulatory site can inhibit the activity of the enzyme





## 5. Product inhibition:

The product of an enzymatic reaction directly binds to the enzyme's active site, competing with the substrate and preventing further catalysis.



• Enzyme is not free



## 6. Induction and repression of enzyme synthesis



→ lag time bet<sup>n</sup> signal and response

- ✓ Induction is a process in which the synthesis of a specific enzyme is activated by expression of its gene.
- ✓ Repression is mechanism by which the synthesis of an enzyme is decreased in response to a specific signal, by inhibiting gene expression.

Basal metabolic Rate (BMR)  
• increase or decrease



## ENZYME PURIFICATION



Total Activity (Decrease)  
% Yield (Decrease)  
Specific activity (Increase)  
Fold purification (Increase)



### Total Activity,(Unit)

- Total activity refers to the overall enzyme activity in a solution at any stage of the purification process.
- Measure of total enzyme in solution

SA

### Specific activity (Unit/mg)

- Activity of an enzyme per milligram of total protein.
- Measure of enzyme purity

### Yield (%)

- Measure of how much of the initial enzyme activity is retained after each purification step.

### Fold Purification


- Increase in purity from the initial extract to various stages of the purification process
- Calculated by dividing the specific activity after each purification step by the specific activity of the initial extract.

$$= \frac{\text{SA after each step}}{\text{SA of crude enzyme}}$$



	Step	Total Activity (Unit)	Total Protein (mg)	Specific Activity (Unit/mg)	Yield (%)	Fold Purification (X)
1	Protein Extract	10000	1000	$\frac{10000}{1000} = 10$	100%	1
2	Salt Precipitation	9000	600	$\frac{9000}{600} = 15$	90%	$\frac{15}{10} = 1.5$
3	Gel Permeation	<del>8000</del> 16000	400	$\frac{8000}{400} = 20$	80%	$\frac{20}{10} = 2$
4	Ion Exchange	7500	250	$\frac{7500}{250} = 30$	75%	$\frac{30}{10} = 3$
5	Affinity Chromatography	7000	100	$\frac{7000}{100} = 70$	70%	$\frac{70}{10} = 7$
		Decrease	Decrease	Increase	Decrease	Increase

- If at some step total activity increase it suggest inhibitor was present prior to this step



- Bond Break
- Oxidation  $R \times^n$
- Energy generate in form ATP, NADH, NADPH.

eg

Glycolysis

Glycogen breakdown

aa oxidation

FA oxidation

- Bond form
- Reduction  $R \times^n$
- Energy is used in form of ATP, NADPH or NADH

eg Gluconeogenesis  
Glycogen biosynthesis  
aa biosynthesis  
FA biosynthesis  
Cholesterol biosynthesis






## GLYCOLYSIS

**Embden-Meyerhof-Parnas (EMP) pathway.**

### Important features:

1. It is a **universal and most conserved pathway**
2. Occurs in **cytosol all cells** of the body- **aerobic/anaerobic**.
3. **Emergency energy**-yielding pathway for cells in the **absence of oxygen**.
4. Major pathway for ATP synthesis in tissues lacking mitochondria, e.g. erythrocytes (RBC), cornea, lens etc.
5. Essential for brain which is dependent on glucose for energy.



# Glycolysis

## Step 1: Hexokinase: First ATP Utilization

**Hexokinase:** *most of tissues*

K<sub>m</sub> for glucose (about 0.1 mM)

Allosteric enzyme

Activator: AMP

Inhibitor: G6P


*ADP  
AMP*

**Glucokinase:** *Liver*

K<sub>m</sub> for glucose (about 10 mM)


MM enzyme

*Low affinity  
for glucose*





## Step 2: Phospho-glucose Isomerase



lysis  
4 + 2


lysis  
3 + 3



## Step 3: Phosphofructokinase: Second ATP Utilization




- major regulatory step
- Pace maker of Glycolysis (slowest)
- committed step of Glycolysis



## Intrinsic Allosteric Regulation of PFK-1: most important

Inhibition: ATP, Citrate, Low pH ( $H^+$ )

Activation: AMP, F6P, F2,6 BP,  $NH_3^+$



PFK-2

- Single bifunctional enzyme
- Kinase & Phosphatase activity


## PFK-2

Inhibitor : Citrate

Activators: F6P, AMP

## B. Splitting phase Step


### 4. Aldolase: Lysis of six carbon compound



Aldolase

Class I : Animals and plants


Class II: Fungi, algae, and some bacteria.



$\uparrow$  DHAP  
Dihydroxy Acetone-3-Phosphate  
 +  
 $\downarrow$  GAP  
Glyceraldehyde-3-phosphate



## Step 5: Triose Phosphate Isomerase



### Inhibitor:

Bromo-hydroxyl-acetone phosphate ✓  
Glycidol phosphate. ✓




## C. Energy generation phase

### Step 6: Oxidation


**Glyceraldehyde-3-Phosphate Dehydrogenase:**  
First "High-Energy" Intermediate Formation

1,3 BP4



Inhibitor: Iodoacetate

E4S  
Irreversible  
inhibitor







## Step 7: Substrate-level phosphorylation

+ 2 ATP

### Phosphoglycerate Kinase: First ATP Generation


✓ Phosphoglycerate kinase reaction is reversible



3 P<sub>4</sub>




## Step 8: Phosphoglycerate Mutase





## Step 9: Enolase:

### Second "High-Energy" Intermediate Formation : PEP




Inhibited by fluoride



## Step 10: Second Substrate-level phosphorylation

+ 2 ATP

### Pyruvate Kinase: Second ATP Generation



Inhibitor: ATP, Alanine




Activator: AMP, F16BP

← Feed forward activation



### Effect of Arsenate


First Substrate level step is bypassed (Net gain 0) ✓



unstable  
Intermediate

## Rapaport-Leubering Cycle of RBC:

First step of ATP synthesis is bypassed and forms 2,3 BPG as intermediate




Bisphosphoglycerate mutase is a bi-functional enzyme with mutase and phosphatase activities





## Fates of Pyruvate under anaerobic condition

### I. Conversion to lactate (Homolactate fermentation):



- Anaerobic condition in muscles
- RBC (aerobic)
- Warburg effect of cancer cells.

Net gain

- only 2 ATP

End Product

- 2 lactate



Hexokinase deficient & Pyruvate Kinase deficient:



NO 2,3 BPG

curve shift  
to left

High 2,3 BPG


curve shift  
to right

%  
Saturation  
↑





## 2. Conversion to ethanol (Alcoholic fermentation):





Requirements:

Net Gain: (2 Pyruvate)

3 Regulatory Kinase & Irreversible Steps:

1 Reversible Kinase :

1 Oxidation step:

2 Energy rich molecules:

2 Substrate level Phosphorylation steps:

2ADP + 2iP + 2NAD + 1Glucose

2ATP + 2 NADH (Aerobic Condition)

2 ATP (Anaerobic Conditions)

1 3 last  
HK (GK), PFK-I, PK

Phosphoglycerate Kinase

Glyceraldehyde 3 P Dehydrogenase

1,3 BPG and PEP

1,3BPG → 3-PG & PEP → Pyruvate

I

II



# Thank you

Learn From India's Best Educators  
India's **No.1** EdTech Company for Graduates  
& Post Graduates Examination.

➡ Download **IFAS** App Now



ANDROID



IOS



WINDOWS





Harshada  
Sharma

**Earned:**  
**₹25,500**

"As is widely recognized, IFAS stands as India's leading institute for CSIR NET examination preparation. My Personal philosophy has always revolved around helping others and I've had the pleasure of introducing the IFAS Online Course to 12 of my classmates and juniors, which in turn allowed me to earn 25,500 rupees. You too can join in and refer your friends, allowing you to earn a 5% rewards based on the fees each of your referred friends Pay"



### Step – 01 Invite Friends

Share your referral code or Link with your friends.

### Step – 02 Earn Cashback

On Every Purchase made using your referral code, you earn 5% instant cashback in your bank account.

### Step – 03 Refer More, Earn More

Keep referring as many friends as you can and keep **Earning** cash as much as you can



Friendship is Sharing, Caring, Giving & Helping Each Other...!